

Quantum holography and multiparticle entanglement using ground-state coherences

G. S. Agarwal¹, P. Lougovski², H. Walther²

¹*Physical Research Laboratory, Navrangpura, Ahmedabad, India*

²*Max-Planck-Institut für Quantenoptik, Hans-Kopfermann-Strasse 1, D-85748 Garching, Germany*

A method of storing and retrieving quantum states of radiation fields using the ground-state coherences is discussed. We demonstrate the generation of multiparticle entangled states starting from atoms prepared in a coherent state. Use is made of the $j=-1/2$ to $+1/2$ atomic transition and its interaction with a far-detuned elliptically polarized field.

PACS numbers: 03.65.Ud 03.67.Mn 42.50.Gy 42.50.Pq

Storage and retrieval of light pulses has been extensively studied both theoretically and experimentally [1, 2, 3, 4, 5, 6, 7]. The light pulses can be stored in atomic coherences and later the atomic coherences can be converted back into the original light pulses [2, 6]. The original work has been extended to storage of pulses of moderate powers [8, 9]. A natural question arises: can one store and retrieve the quantum state of the field? This is the question addressed in this letter. It is shown how coherently prepared atomic systems interacting with far-detuned elliptical fields can serve the purpose. It is also demonstrated how the atomic coherences can be used for generating a variety of nonclassical states, including multiatom entangled states. Our model consists of N identical two-level atoms, with hyperfine structure splitting as depicted in Fig.1, coupled to two mutually orthogonal modes of a quantized electromagnetic field described by creation (annihilation) operators $a_{-}^{\dagger}, a_{+}^{\dagger}$ (a_{-}, a_{+}) with the ground and excited-state coupling strengths Ω_{+} and Ω_{-} , respectively. The Hamiltonian for the atom-field system reads

$$\begin{aligned} H = & \hbar \frac{\omega_1}{2} \sum_{j=1}^N (|e_{-}\rangle\langle e_{-}| - |g_{+}\rangle\langle g_{+}|)_j \\ & + \hbar \frac{\omega_2}{2} \sum_{j=1}^N (|e_{+}\rangle\langle e_{+}| - |g_{-}\rangle\langle g_{-}|)_j \\ & + \hbar \omega_f (a_{+}^{\dagger} a_{+} + a_{-}^{\dagger} a_{-}) \\ & + \hbar (\Omega_{-} a_{-} \sum_{j=1}^N |e_{-}\rangle\langle g_{+}|_j + h.c.) \\ & + \hbar (\Omega_{+} a_{+} \sum_{j=1}^N |e_{+}\rangle\langle g_{-}|_j + h.c.), \end{aligned} \quad (1)$$

where ω_1 and ω_2 are the frequencies of the transitions $|g_{+}\rangle \leftrightarrow |e_{-}\rangle$ and $|g_{-}\rangle \leftrightarrow |e_{+}\rangle$, respectively, and ω_f is the field frequency. In the dispersive limit of the atom-field coupling, i.e. when $\omega_1 - \omega_f = \omega_2 - \omega_f = \Delta \gg$

$\{| \Omega_{+} |, | \Omega_{-} | \}$, one can reduce the Hamiltonian Eq.(1) to

$$\begin{aligned} H_1 = & \hbar \frac{|\Omega_{-}|^2}{\Delta} a_{-}^{\dagger} a_{-} \sum_{j=1}^N (|e_{-}\rangle\langle e_{-}| - |g_{+}\rangle\langle g_{+}|)_j \\ & + \hbar \frac{|\Omega_{+}|^2}{\Delta} a_{+}^{\dagger} a_{+} \sum_{j=1}^N (|e_{+}\rangle\langle e_{+}| - |g_{-}\rangle\langle g_{-}|)_j \\ & + \frac{2\hbar}{\Delta} (|\Omega_{-}|^2 \sum_{j=1}^N |e_{-}\rangle\langle e_{-}|_j + |\Omega_{+}|^2 \sum_{j=1}^N |e_{+}\rangle\langle e_{+}|_j). \end{aligned} \quad (2)$$

The effective Hamiltonian Eq.(2), in contrast to the original Eq.(1), does not couple ground states $\{|g_{-}\rangle_j, |g_{+}\rangle_j\}$, $j = 1, N$ to the excited ones. Therefore, if one starts from the initial atomic ground state, the only relevant contribution from the Hamiltonian Eq.(2) is

$$\begin{aligned} H_r = & -\hbar \frac{|\Omega_{-}|^2}{\Delta} a_{-}^{\dagger} a_{-} \sum_{j=1}^N (|g_{+}\rangle\langle g_{+}|)_j \\ & - \hbar \frac{|\Omega_{+}|^2}{\Delta} a_{+}^{\dagger} a_{+} \sum_{j=1}^N (|g_{-}\rangle\langle g_{-}|)_j. \end{aligned} \quad (3)$$

Introducing the collective atomic operator

$$\hat{R}_z = \frac{1}{2} \sum_{j=1}^N (|g_{+}\rangle\langle g_{+}| - |g_{-}\rangle\langle g_{-}|)_j, \quad (4)$$

one can express the operators $\sum_{j=1}^N (|g_{+}\rangle\langle g_{+}|)_j$ and $\sum_{j=1}^N (|g_{-}\rangle\langle g_{-}|)_j$ in terms of $\hat{R}_z$ as follows:

$$\frac{1}{2} \pm \hat{R}_z = \sum_{j=1}^N (|g_{\pm}\rangle\langle g_{\pm}|)_j.$$

For the sake of simplicity hereafter we suppose that the two field modes have identical coupling strengths, i.e. $|\Omega_{+}| = |\Omega_{-}| = \Omega$. Finally, the effective Hamiltonian Eq.(2) will read

$$H' = \frac{\hbar \Omega^2}{\Delta} \hat{N}_z \hat{R}_z, \quad (5)$$

where we have introduced the following field operator:

$$\hat{N}_z = a_+^\dagger a_+ - a_-^\dagger a_-.$$

Apart from the $\hat{R}_z$ operator, we construct the lowering $\hat{R}_-$ and raising $\hat{R}_+$ operators,

$$\hat{R}_- = \sum_{j=1}^N (|g_-\rangle\langle g_+|)_j, \quad (6)$$

$$\hat{R}_+ = \sum_{j=1}^N (|g_+\rangle\langle g_-|)_j. \quad (7)$$

The operators $\hat{R}_\pm$ and $\hat{R}_z$ fulfill the following commutation relations:

$$[\hat{R}_-, \hat{R}_+] = -2\hat{R}_z, [\hat{R}_z, \hat{R}_\pm] = \pm\hat{R}_\pm, \quad (8)$$

and therefore describe an effective spin $J = \frac{N}{2}$ system, where N is the number of atoms in the sample. As next step we define the eigenvectors of the z component of the total spin operator $\hat{R}_z$ as follows:

$$\hat{R}_z|m\rangle = \hbar m|m\rangle, J = \frac{N}{2}, m = -J, -J+1, \dots, J-1, J. \quad (9)$$

Action of the lowering (raising) operators $\hat{R}_-$ ($\hat{R}_+$) on the eigenvectors of $\hat{R}_z$ can be calculated in a straightforward manner:

$$\hat{R}_-|m\rangle = \hbar\sqrt{(J+m)(J-m+1)}|m-1\rangle, \quad (10)$$

$$\hat{R}_+|m\rangle = \hbar\sqrt{(J-m)(J+m+1)}|m+1\rangle. \quad (11)$$

The set of eigenvectors $|m\rangle, m = \overline{-J, J}$ of the operator $\hat{R}_z$ is trivially connected to the previously introduced set $\{|g_-\rangle_j, |g_+\rangle_j\}, j = \overline{1, N}$, which denotes the hyperfine structure of the ground states of atoms. For example $|-J\rangle = \prod_{k=1}^N |g_-\rangle_k$, and consequently we will call the state $|-J\rangle$ the ground state. It is instructive to introduce the following field operators,

$$\hat{N}_- = a_-^\dagger a_+, \hat{N}_+ = a_+^\dagger a_-, \quad (12)$$

and compare the algebraic properties of the two sets of operators, $\hat{N}_z, \hat{N}_-, \hat{N}_+$ and $\hat{R}_z, \hat{R}_-, \hat{R}_+$. Operators $\hat{N}_z, \hat{N}_-$ and $\hat{N}_+$ satisfy the same commutation relations as in Eq.(8), i.e.

$$[\hat{N}_-, \hat{N}_+] = -2\hat{N}_z, [\hat{N}_z, \hat{N}_\pm] = \pm\hat{N}_\pm. \quad (13)$$

Therefore, coming back to the Hamiltonian Eq.(5), one can say that there is effectively a spin-spin interaction along the z-axis, where one of the spins is the collective spin of the atomic sample and the other one is some effective spin which corresponds to two orthogonally polarized modes of the cavity field. As a result, the Hamiltonian Eq.(5) is very suitable for spin quantum nondemolition




FIG. 1: Internal atomic structure

measurement (QND) as well as being a good candidate for the generation of spin-squeezed atomic samples [10].

We now demonstrate important applications of the Hamiltonian Eq.(5) to create mesoscopic superposition states [11], multiatom entangled states and to store and retrieve quantum states.

Mesoscopic superposition of states: Let us consider the case where the atomic sample has been initially prepared in some coherent atomic state [12]

$$\begin{aligned} |\theta, \phi\rangle &= R_{\theta, \phi}|-J\rangle \\ &= \sum_{m=-J}^J \binom{2J}{J+m}^{\frac{1}{2}} \left[\cos\left(\frac{\theta}{2}\right) \right]^{J-m} \left[\sin\left(\frac{\theta}{2}\right) e^{-i\phi} \right]^{J+m} |m\rangle. \end{aligned} \quad (14)$$

Here $R_{\theta, \phi}$ is the rotation operator which rotates the initial ground state $|-J\rangle$ by an angle θ around an axis $\hat{m} = (\sin(\phi), -\cos(\phi), 0)$. From the practical point of view the coherent atomic state in Eq.(14) can be prepared in a number of ways; for example, by applying a homogeneous magnetic field along the $\hat{m}$ axis to the ground state of the atomic sample $|-J\rangle$. The other interesting and popular possibility is to use short pulses and Raman transitions, a technique that has been extensively used by, for example, Wineland and co-workers [13]. We consider interaction of the coherently prepared atomic sample with an elliptically polarized field which we represent by the state $|\psi_f\rangle = \alpha|1_+, 0_-\rangle + \beta|0_+, 1_-\rangle$, ($|\alpha|^2 + |\beta|^2 = 1$). Note that $\alpha = 0$ or $\beta = 0$ corresponds to a circularly polarized field. A calculation by means of the interaction Hamiltonian Eq.(5) leads to the following atom-field state

$$\begin{aligned} |\Psi_{at-f}\rangle &= \exp(-i\phi_0 t \hat{N}_z \hat{R}_z) |\theta, \phi\rangle |\psi_f\rangle \\ &= \alpha |\theta, \phi + \phi_0 t\rangle |1_+, 0_-\rangle e^{i\phi_0 t J} \\ &\quad + \beta |\theta, \phi - \phi_0 t\rangle |0_+, 1_-\rangle e^{-i\phi_0 t J}. \end{aligned} \quad (15)$$

Here $\phi_0 = \Omega^2/\Delta$ and t is the interaction time. The atom-field state Eq.(16) describes a situation when atoms and the cavity field get entangled. Detection of the field polarization in the direction of, say, x leads to an atomic

state

$$|\Psi\rangle_{cat} = \alpha e^{i\phi_0 t J} |\theta, \phi + \phi_0 t\rangle + \beta e^{-i\phi_0 t J} |\theta, \phi - \phi_0 t\rangle. \quad (16)$$

This corresponds to a mesoscopic superposition of coherent states for the atomic system. This is the case as long as the two coherent states in the above equation are not orthogonal, i.e.

$$\langle \theta, \phi + \phi_0 t | \theta, \phi - \phi_0 t \rangle = \exp(2iJ\phi_0 t) [\cos(\phi_0 t) - i \cos(\theta) \sin(\phi_0 t)]^{2J}. \quad (17)$$

is not zero. From Eq.(17) follows that for a particular choice of parameters (for example, $\phi_0 t = \frac{\pi}{2}$ and $\theta = \frac{\pi}{2}$, $\forall \phi$) the overlap between coherent atomic states is zero and hence, the atom-field state in Eq.(16) becomes a mixed state. We have thus shown how the dispersive interactions on ground-state atoms can produce long-lived mesoscopic superpositions. We note that dispersive interaction has been considered in the context of two-level atoms to produce both atomic and field cat states [14, 15, 16, 17].

Generation of a multiatom entangled state: It is next shown that the above mesoscopic superposition is equivalent to a multiatom entangled state. For this purpose we express the coherent state of Eq. (14) as a product state over all the atoms:

$$|\theta, \phi\rangle = \prod_{j=1, N} |\theta, \phi\rangle_j \\ |\theta, \phi\rangle_j = \sin(\theta/2) e^{-i\phi} |g_+\rangle_j + \cos(\theta/2) |g_-\rangle_j, \quad (18)$$

and hence the mesoscopic superposition (16) can be expressed as

$$|\psi\rangle_{cat} = \alpha \prod_{j=1, N} |\theta, \phi + \phi_0 t\rangle_j e^{i\phi_0 t J} + \beta e^{-i\phi_0 t J} \prod_{j=1, N} |\theta, \phi - \phi_0 t\rangle_j. \quad (19)$$

This is a multi-atom entangled state. We further note that $\langle \theta, \phi - \phi_0 t | \theta, \phi + \phi_0 t \rangle = 0$ if $\phi_0 t = \pi/2$, $\theta = \pi/2$. Thus the two parts of (19) become orthogonal for this choice of parameters. In this case (19) becomes an example of GHZ states [18]. It has thus been shown how long-lived GHZ states can be produced by using ground-state atomic coherences.

Interaction-free evolution of field states: It is clear from Eq.(15) that the probability of detecting the field state unchanged after the coupling to atoms strongly depends on the atom-field interaction time. If we choose $\phi_0 t = \pi$ and make the number of atoms in the sample N even, the cavity field after interaction is in the same state as the initial cavity field. This is reminiscent of the trapping states in a micromaser [19] where for certain field states the atoms leave the cavity in the original states. To verify the above statement we consider the evolution operator $\exp\{-i\pi \hat{N}_z \hat{R}_z\}$ for $\phi_0 t = \pi$. Since

$\hat{N}_z$ has the eigenvalues ± 1 and for even N , $\hat{R}_z$ has an integer eigenvalue and the above evolution operator is the same for the two eigenvalues of $\hat{N}_z$. Thus the field would revert to the original state. For odd N , the eigenvalues of $(\frac{N}{2} + \hat{R}_z)$ are integers and then the two evolution operators $\exp\{\mp i\pi \hat{R}_z\}$ will differ by an overall phase factor π ($\exp\{2i\pi N/2\}$). In this case the state of the output field will be $\alpha|1_+, 0_-\rangle - \beta|0_+, 1_-\rangle$.

Quantum holography : It is next demonstrated how our system can be used to store the quantum state of the field via atomic coherence. This is the quantum counterpart of holography in which the object information is stored in a hologram using a reference field. The hologram when irradiated by a coherent field, recovers the information on the object. Clearly Eq.(15) already shows how the information about the field state is stored in atomic ground-state coherence. Note that this is long-lived storage since we use the ground states of the atomic systems. It is now shown how this quantum hologram can be read to retrieve the information on the field. One possibility is to measure the atomic population in the state $|g_-\rangle$. This is easily done by applying a laser pulse polarized appropriately so as to move the population from $|g_-\rangle$ (and not from $|g_+\rangle$) to an excited state and to measure the subsequent fluorescence. This measurement will project the state (15) on to a state of the field given by

$$|\psi_f\rangle_{-J} \rightarrow \left(\cos \frac{\theta}{2}\right)^{2J} \times (\alpha|1_+, 0_-\rangle e^{i\phi_0 t J} + \beta|0_+, 1_-\rangle e^{-i\phi_0 t J}). \quad (20)$$

Thus the stored state of the field is recovered apart from a relative phase factor $2J\phi_0 t$. Note that this phase factor is known a priori from the preparation process of the hologram.

Generation of cat states for the radiation field: Finally, it is shown how the interaction (5) is quite suited to production of cat states for the radiation field. This possibility has already been extensively discussed by Haroche and co-workers[17] using Rydberg atoms. Here we use ground-state coherence. Consider, for example, a single atom $J = 1/2$ interacting with a field in coherent state $|\alpha, \beta\rangle$. Obviously, the state of the combined system at time t will be

$$|\psi(t)\rangle \equiv \exp\{-i\phi_0 t \hat{N}_z \hat{R}_z\} |\alpha, \beta\rangle |\theta, \phi\rangle \\ = \cos \frac{\theta}{2} |g_-\rangle |\alpha e^{i\phi_0 t/2}, \beta e^{-i\phi_0 t/2}\rangle \\ + e^{-i\phi} \sin \frac{\theta}{2} |g_+\rangle |\alpha e^{-i\phi_0 t/2}, \beta e^{i\phi_0 t/2}\rangle. \quad (21)$$

Clearly, measurement of the atomic population in a state different from $|g_\pm\rangle$ produces a cat state of the radiation field.

In conclusion, it has been shown how ground-state coherence can be utilized for quantum holography and for

generation of long-lived mesoscopic superpositions and multi particle GHZ states. The latter states possess very interesting nonclassical properties.

-
- [1] C. Liu, Z. Dutton, C.H. Behroozi, and L.V. Hau, *Nature* (London) **409**, 490 (2001).
 - [2] M. Fleischhauer and M.D. Lukin, *Phys. Rev. Lett.* **84**, 5094 (2000); M. Fleischhauer and M.D. Lukin, *Phys. Rev. A* **65**, 022314 (2002); D.F. Phillips, A. Fleischhauer, A. Mair, R.L. Walsworth, and M.D. Lukin, *Phys. Rev. Lett.* **86**, 783 (2001); J. R. Czeszegi and R. Grobe, *Phys. Rev. Lett.* **79**, 3162 (1997).
 - [3] A.S. Zibrov, A.B. Matsko, O. Kocharovskaya, Y.V. Rostovtsev, G.R. Welch, and M.O. Scully, *Phys. Rev. Lett.* **88**, 103601 (2002).
 - [4] A.B. Matsko, Y.V. Rostovtsev, O. Kocharovskaya, A.S. Zibrov, and M.O. Scully, *Phys. Rev. A* **64**, 043809 (2001).
 - [5] A. Kuzmich, W. P. Bowen, A. D. Boozer, A. Boca, C. W. Chou, L.-M. Duan, H. J. Kimble, *Nature* **423**, 731 (2003).
 - [6] M. G. Payne and L. Deng, *Phys. Rev. Lett.* **91**, 123602 (2003).
 - [7] C. H. van der Wal, M. D. Eisaman, A. Andr , R. L. Walsworth, D. F. Phillips, A. S. Zibrov, and M. D. Lukin *Science* **301** 196 (2003).
 - [8] T. N. Dey and G. S. Agarwal, *Phys. Rev. A* **67**, 0433813 (2003).
 - [9] V. G. Arkhipkin and I. V. Timofeev, *JETP Letters* **76**, 66 (2002).
 - [10] L.M.Duan et al., *Phys.Rev.Lett.* **85**, 5643 (2000), A.Kuzmich and E. S. Polzik, *Phys. Rev. Lett.* **85**, 5639 (2000), A.Kuzmich, L.Mandel, and N.P.Bigelow, *Phys. Rev. Lett.* **85**, 1594 (2000), S.Massar and E.S.Polzik, *Phys. Rev. Lett.* **91**, 060401 (2003).
 - [11] E.Schr dinger, *Naturwissenschaften* **23**, pp. 807-812; 823-828; 844-849 (1935).
 - [12] F.T.Arecchi, E.Courtens, R.Gilmore, H. Thomas, *Phys.Rev.A* **6**, 2211 (1972).
 - [13] C. Monroe, D. M. Meekhof, B. E. King, and D. J. Wineland, *Science* **272**, 1131 (1996).
 - [14] G. S. Agarwal, R. R. Puri, and R. P. Singh, *Phys. Rev. A* **56**, 2249 (1997).
 - [15] C. C. Gerry, and R. Grobe, *Phys. Rev. A* **57**, 2247 (1998), C. C. Gerry *ibid* **54**, 2529(1996).
 - [16] S. B. Zeng, *Phys. Rev. Lett.* **87**, 230404 (2001).
 - [17] M. Brune, E. Hagley, J. Dreyer, X. Maitre, A. Maali, C. Wunderlich, J. M. Raimond, and S. Haroche, *Phys. Rev. Lett.* **77**, 4887 (1996).
 - [18] D.M. Greenberger, M.A. Horne, and A. Zeilinger, *Phys. Today* **46**, No.8, 22 (1993).
 - [19] P. Meystre, G. Rempe, and H. Walther, *Opt. Lett.* **13**, 1078 (1988).